

On the Stability of J^* -Homomorphisms ^{*}

Choonkil Baak and Mohammad Sal Moslehian

Abstract

The main purpose of this paper is to prove the generalized Hyers-Ulam-Rassias stability of J^* -homomorphisms between J^* -algebras.

1 Introduction

In 1940, S. M. Ulam posed the following problem concerning the stability of group homomorphisms [13]:

Given a group G_1 , a metric group (G_2, d) and a positive number ϵ , does there exist a $\delta > 0$ such that if a function $f : G_1 \rightarrow G_2$ satisfies the inequality $d(f(xy), f(x)f(y)) < \delta$ for all $x, y \in G_1$ then there exists a homomorphism $T : G_1 \rightarrow G_2$ such that $d(f(x), T(x)) < \epsilon$ for all $x \in G_1$. If this problem has a solution, we say that the homomorphisms from G_1 to G_2 are stable or the functional equation $f(xy) = f(x)f(y)$ is stable.

In 1941, D. H. Hyers gave a partial solution of Ulam's problem in the context of Banach spaces as the following [7]:

Suppose that E_1, E_2 are Banach spaces and $f : E_1 \rightarrow E_2$ satisfies the following condition: there is $\epsilon > 0$ such that $\|f(x+y) - f(x) - f(y)\| < \epsilon$ for all $x, y \in E_1$. Then there is an additive mapping $T : E_1 \rightarrow E_2$ such that $\|f(x) - T(x)\| < \epsilon$ for all $x \in E_1$.

^{*}2000 *Mathematics Subject Classification*. Primary 39B82; Secondary 47B48, 46L05, 39B52, 46K99, 16Wxx.

Keywords and phrases. Hyers-Ulam-Rassias stability, J^* -algebra, J^* -homomorphism, C^* -algebra.

The first author was supported by Korea Research Foundation Grant KRF-2004-041-C00023.

Now, assume that E_1 and E_2 are real normed spaces with E_2 complete, $f : E_1 \rightarrow E_2$ is a mapping such that $f(tx)$ is continuous in $t \in \mathbb{R}$ for each fixed $x \in E_1$, and let there exist $\varepsilon \geq 0$ and $p \in [0, \infty) - \{1\}$ such that

$$\|f(x+y) - f(x) - f(y)\| \leq \varepsilon(\|x\|^p + \|y\|^p),$$

for all $x, y \in E_1$. It was shown by Th. M. Rassias [11] for $p \in [0, 1)$ and Z. Gajda [2] for $p > 1$ that there exists a unique linear map $T : E_1 \rightarrow E_2$ such that

$$\|f(x) - T(x)\| \leq \frac{2\varepsilon}{|2^p - 2|} \|x\|^p,$$

for all $x \in E_1$. Using Hyers' method, indeed, $T(x)$ is defined by $\lim_{n \rightarrow \infty} 2^{-n} f(2^n x)$ if $0 \leq p < 1$, and $\lim_{n \rightarrow \infty} 2^n f(2^{-n} x)$ if $p > 1$. This phenomenon is called the Hyers–Ulam–Rassias stability. It is shown that there is no analogue of Rassias result for $p = 1$ (see [12]).

In 1992, a generalization of Rassias' theorem was obtained by Găvruta as follows [3]:

Suppose $(G, +)$ is an abelian group, E is a Banach space and the so-called admissible control function $\varphi : G \times G \rightarrow [0, \infty)$ satisfies

$$\tilde{\varphi}(x, y) := \frac{1}{2} \sum_{n=0}^{\infty} 2^{-n} \varphi(2^n x, 2^n y) < \infty,$$

for all $x, y \in G$. If $f : G \rightarrow E$ is a mapping with

$$\|f(x+y) - f(x) - f(y)\| \leq \varphi(x, y),$$

for all $x, y \in G$, then there exists a unique mapping $T : G \rightarrow E$ such that $T(x+y) = T(x) + T(y)$ and $\|f(x) - T(x)\| \leq \tilde{\varphi}(x, x)$ for all $x, y \in G$.

During the last decades several stability problems of functional equations have been investigated in spirit of Hyers–Ulam–Rassias, see [1].

By a J^* -algebra we mean a closed subspace A of a C^* -algebra such that $xx^*x \in A$ whenever $x \in A$. Many familiar spaces are J^* -algebras [4]. For example, (i) every Cartan factor of type I , i.e., the space of all bounded operators $B(H, K)$ between Hilbert spaces H and K ; (ii) every Cartan factor of type IV , i.e., a closed $*$ -subspace A of $B(H)$ in which the square of each operator in A is a scalar multiple of the identity operator on H ; (iii) every JC^* -algebra; (iv) every ternary algebra of operators [5].

A J^* -homomorphism between J^* -algebras A and B is defined to be a linear mapping $L : A \rightarrow B$ such that $T(xx^*x) = Tx(Tx)^*Tx$ for all $x \in A$. In particular, every $*$ -homomorphism between C^* -algebras is a J^* -homomorphism.

In [9], the first author establishes the stability of $*$ -homomorphisms of a C^* -algebra. Using Rassias' technique [11] and some similar methods as in [10], we generalize some of the results for J^* -homomorphisms.

Throughout this paper, A and B denote J^* -algebras.

2 Main results

Our main purpose is to prove the generalized Hyers–Ulam–Rassias stability of J^* -homomorphisms.

Proposition 2.1. *Let $r > 1$, and let $T : A \rightarrow A$ be a mapping satisfying $T(rx) = rT(x)$ for all $x \in A$ and let there exist a function $\varphi : A \times A \times A \rightarrow [0, \infty)$ such that*

$$\lim_{n \rightarrow \infty} r^{-n} \varphi(r^n x, r^n y, r^n z) = 0,$$

$$\|T(\lambda x + y + zz^*z) - \lambda T(x) - T(y) - T(z)T(z)^*T(z)\| \leq \varphi(x, y, z), \quad (1)$$

for all $\lambda \in \mathbf{C}$ and all $x, y, z \in A$. Then T is a J^* -homomorphism.

Proof. $T(0) = 0$ since $T(0) = rT(0)$. Put $x = y = 0$ in (1). Then

$$\begin{aligned} \|T(zz^*z) - T(z)T(z)^*T(z)\| &= \frac{1}{r^{3n}} \|T(r^n z(r^n z)^*r^n z) - T(r^n z)T(r^n z)^*T(r^n z)\| \\ &\leq \frac{1}{r^{3n}} \varphi(r^n z, r^n z, r^n z) \leq \frac{1}{r^n} \varphi(r^n z, r^n z, r^n z), \end{aligned}$$

for all $z \in A$. The right side tends to zero as $n \rightarrow \infty$. So

$$T(zz^*z) = T(z)T(z)^*T(z),$$

for all $z \in A$.

Similarly, one can show that

$$T(\lambda x + y) = \lambda T(x) + T(y),$$

for all $x, y \in A$. $\square$

Theorem 2.2. Suppose $h : A \rightarrow B$ is a mapping with $h(0) = 0$ for which there exists a function $\varphi : A \times A \times A \rightarrow [0, \infty)$ such that

$$\tilde{\varphi}(x, y, z) := \frac{1}{2} \sum_{n=0}^{\infty} 2^{-n} \varphi(2^n x, 2^n y, 2^n z) < \infty,$$

$$\|h(\mu x + \mu y + zz^*z) - \mu h(x) - \mu h(y) - h(z)h(z)^*h(z)\| \leq \varphi(x, y, z), \quad (2)$$

for all $\mu \in S^1 = \{\lambda \in \mathbf{C} : |\lambda| = 1\}$ and all $x, y, z \in A$. Then there exists a unique J^* -homomorphism $T : A \rightarrow B$ such that

$$\|h(x) - T(x)\| \leq \tilde{\varphi}(x, x, 0),$$

for all $x \in A$.

Proof. In (2), assume that $z = 0$ and $\mu = 1$. Then the Găvruta theorem implies that there is a unique additive mapping $T : A \rightarrow B$ given by

$$T(x) = \lim_{n \rightarrow \infty} \frac{h(2^n x)}{2^n},$$

for all $x \in A$. By (2),

$$\|h(2^n \mu x) - 2\mu h(2^{n-1}x)\| \leq \varphi(2^{n-1}x, 2^{n-1}x, 0),$$

for all $x \in A$ and all $\mu \in S^1$. Then

$$\|\mu h(2^n x) - 2\mu h(2^{n-1}x)\| \leq |\mu| \cdot \|h(2^n x) - 2h(2^{n-1}x)\| \leq \varphi(2^{n-1}x, 2^{n-1}x, 0),$$

for all $\mu \in S^1$ and all $x \in A$. So

$$\begin{aligned} \|2^{-n}h(2^n \mu x) - \mu 2^{-n}h(2^n x)\| &\leq 2^{-n}\|h(2^n \mu x) - 2\mu h(2^{n-1}x)\| + 2^{-n}\|2\mu h(2^{n-1}x) - \mu h(2^n x)\| \\ &\leq 2^{-n+1}\varphi(2^{n-1}x, 2^{n-1}x, 0), \end{aligned}$$

for all $\mu \in S^1$ and all $x \in A$. Since the right side tends to zero as $n \rightarrow \infty$, we have

$$T(\mu x) = \lim_{n \rightarrow \infty} \frac{h(2^n \mu x)}{2^n} = \lim_{n \rightarrow \infty} \frac{\mu h(2^n x)}{2^n} = \mu T(x),$$

for all $\mu \in S^1$ and all $x \in A$. Obviously, $T(0x) = 0 = 0T(x)$.

Next, let $\lambda \in \mathbf{C} (\lambda \neq 0)$ and let M be a natural number greater than $4|\lambda|$. Then $|\frac{\lambda}{M}| < \frac{1}{4} < 1 - \frac{2}{3} = 1/3$. By Theorem 1 of [8], there exist three numbers $\mu_1, \mu_2, \mu_3 \in S^1$

such that $3\frac{\lambda}{M} = \mu_1 + \mu_2 + \mu_3$. By the additivity of T we get $T(\frac{1}{3}x) = \frac{1}{3}T(x)$ for all $x \in A$. Therefore,

$$\begin{aligned} T(\lambda x) &= T\left(\frac{M}{3} \cdot 3 \cdot \frac{\lambda}{M}x\right) = MT\left(\frac{1}{3} \cdot 3 \cdot \frac{\lambda}{M}x\right) = \frac{M}{3}T\left(3 \cdot \frac{\lambda}{M}x\right) \\ &= \frac{M}{3}T(\mu_1 x + \mu_2 x + \mu_3 x) = \frac{M}{3}(T(\mu_1 x) + T(\mu_2 x) + T(\mu_3 x)) \\ &= \frac{M}{3}(\mu_1 + \mu_2 + \mu_3)T(x) = \frac{M}{3} \cdot 3 \cdot \frac{\lambda}{M} = \lambda T(x), \end{aligned}$$

for all $x \in A$. So that T is $\mathbf{C}$ -linear.

Put $x = y = 0$ and replace z by $2^n z$ in (2). Then

$$\frac{1}{2^{3n}} \|h(2^{3n} z z^* z) - h(2^n z)h(2^n z)^*h(2^n z)\| \leq \frac{1}{2^{3n}} \varphi(0, 0, 2^n z) \leq \frac{1}{2^n} \varphi(0, 0, 2^n z),$$

for all $z \in A$. Hence

$$\begin{aligned} T(z z^* z) &= \lim_{n \rightarrow \infty} \frac{h(2^{3n} z z^* z)}{2^{3n}} = \lim_{n \rightarrow \infty} \frac{h(2^n z)}{2^n} \frac{h(2^n z)^*}{2^n} \frac{h(2^n z)}{2^n} \\ &= T(z)T(z)^*T(z), \end{aligned}$$

for all $z \in A$. It follows that T is a J^* -homomorphism satisfying the required inequality. $\square$

Example 2.3. Let $H : A \rightarrow A$ be a J^* -homomorphism, and let $h : A \rightarrow A$ be defined by $h(x) = \begin{cases} H(x) & \|x\| < 1 \\ 0 & \|x\| \geq 1 \end{cases}$ and $\varphi(x, y, z) = 4$.

Then

$$\tilde{\varphi}(x, y, z) = \frac{1}{2} \sum_{n=0}^{\infty} 2^{-n} \cdot 4 = 4,$$

and

$$\|h(\mu x + \mu y + z z^* z) - \mu h(x) - \mu h(y) - h(z)h(z)^*h(z)\| \leq 4 = \varphi(x, y, z),$$

for all $\mu \in S^1$ and all $x, y, z \in A$.

Since in a C^* -algebra

$$\|xx^*x\| = \|xx^*xx^*xx^*\|^{\frac{1}{2}} = \|(xx^*)^3\|^{\frac{1}{2}} = \|xx^*\|^{\frac{3}{2}} = \|x\|^{2 \cdot \frac{3}{2}} = \|x\|^3,$$

and every J^* -homomorphism is norm decreasing [6], we conclude that $h(xx^*x) = h(x)h(x^*)h(x)$.

Note also that h is not linear. Further, $T(0) = \lim_{n \rightarrow \infty} \frac{h(0)}{2^n} = 0$ and for $x \neq 0$ we have

$$T(x) = \lim_{n \rightarrow \infty} \frac{h(2^n x)}{2^n} = \lim_{n \rightarrow \infty} \frac{0}{2^n} = 0,$$

since for sufficiently large n , $\|2^n x\| \geq 1$. Thus T is identically zero and

$$\|h(x) - T(x)\| \leq \tilde{\varphi}(x, x, 0) = 4,$$

for all $x \in A$.

Corollary 2.4. *Suppose that $h : A \rightarrow B$ is a mapping with $h(0) = 0$ for which there exist constants $\alpha \geq 0$ and $p \in [0, 1)$ such that*

$$\|h(\mu x + \mu y + zz^*z) - \mu h(x) - \mu h(y) - h(z)h(z)^*h(z)\| \leq \alpha(\|x\|^p + \|y\|^p + \|z\|^p),$$

for all $\mu \in S^1$ and all $x, y, z \in A$. Then there is a unique J^* -homomorphism $T : A \rightarrow B$ such that

$$\|h(x) - T(x)\| \leq \frac{\alpha}{1 - 2^{p-1}} \|x\|^p,$$

for all $x \in A$.

Proof. Put $\varphi(x, y, z) = \alpha(\|x\|^p + \|y\|^p + \|z\|^p)$ in Theorem 2.2. $\square$

Theorem 2.5. *Suppose that $h : A \rightarrow B$ is a mapping with $h(0) = 0$ for which there exists a function $\varphi : A \times A \times A \rightarrow [0, \infty)$ such that*

$$\tilde{\varphi}(x, y, z) := \frac{1}{2} \sum_{n=0}^{\infty} 2^{-n} \varphi(2^n x, 2^n y, 2^n z) < \infty,$$

$$\|h(\mu x + \mu y + zz^*z) - \mu h(x) - \mu h(y) - h(z)h(z)^*h(z)\| \leq \varphi(x, y, z), \quad (3)$$

for $\mu = 1, i$ and all $x, y, z \in A$. If for each fixed $x \in A$ the function $t \mapsto h(tx)$ is continuous on $\mathbf{R}$, then there exists a unique J^* -homomorphism $T : A \rightarrow B$ such that

$$\|h(x) - T(x)\| \leq \tilde{\varphi}(x, x, 0),$$

for all $x \in A$.

Proof. Put $z = 0$ and $\mu = 1$ in (3). It follows from the Găvruta theorem that there exists a unique additive mapping $T : A \rightarrow B$ given by

$$T(x) = \lim_{n \rightarrow \infty} \frac{h(2^n x)}{2^n},$$

for all $x \in A$.

By the same reasoning as in the proof of the main theorem of [11], the mapping T is $\mathbf{R}$ -linear.

Assume $y = z = 0$ and $\mu = i$. It follows from (3) that

$$\|h(ix) - ih(x)\| \leq \varphi(x, 0, 0),$$

for all $x \in A$. Hence

$$\frac{1}{2^n} \|h(2^n ix) - ih(2^n x)\| \leq \varphi(2^n x, 0, 0),$$

for all $x \in A$. The right side tends to zero as $n \rightarrow \infty$, so

$$T(ix) = \lim_{n \rightarrow \infty} \frac{h(2^n ix)}{2^n} = \lim_{n \rightarrow \infty} \frac{ih(2^n x)}{2^n} = iT(x),$$

for all $x \in A$. For every $\lambda \in \mathbf{C}$, $\lambda = s + it$ in which $s, t \in \mathbf{R}$ we have

$$\begin{aligned} T(\lambda x) &= T(sx + itx) = sT(x) + tT(ix) = sT(x) + itT(x) = (s + it)T(x) \\ &= \lambda T(x), \end{aligned}$$

for all $x \in A$. Thus T is $\mathbf{C}$ -linear. $\square$

References

- [1] S. Czerwik, Stability of Functional Equations of Ulam–Hyers–Rassias Type, Hadronic Press, 2003.
- [2] Z. Gajda, On stability of additive mappings, Int. J. Math. Math. Sci. 14 (1991), 431–434.
- [3] P. Găvruta, A generalization of the Hyers–Ulam–Rassias stability of approximately additive mappings, J. Math. Anal. Appl. 184 (1994), 431–436.
- [4] L.A. Harris, Bounded Symmetric Homogeneous Domains in Infinite-Dimensional Spaces, Lecture Notes in Mathematics 364, Springer, Berlin, 1974.
- [5] L.A. Harris, Operator Siegel domains”, Proc. Roy. Soc. Edinburgh Sect. A. 79 (1977), 137–156.
- [6] L.A. Harris, A generalization of C^* -algebras, Proc. London Math. Soc. 42 (1981), 331–361.

- [7] D.H. Hyers, On the stability of the linear functional equation, Proc. Nat. Acad. Sci. U.S.A. 27 (1941), 222–224.
- [8] R.V. Kadison and G.K. Pedersen, Means and convex combinations of unitary operators, Math. Scan. 57 (1985), 249–266.
- [9] C. Park, On an approximate automorphism on a C^* -algebra, Proc. Amer. Math. Soc. 132 (2004), 1739–1745.
- [10] C. Park, Lie $*$ -homomorphisms between Lie C^* -algebras and Lie $*$ -derivations on Lie C^* -algebras, J. Math. Anal. Appl. 293 (2004), 419–434.
- [11] Th.M. Rassias, On the stability of the linear mapping in Banach spaces, Proc. Amer. Math. Soc. 72 (1978), 297–300.
- [12] Th.M. Rassias and P. Šemrl, On the behavior of mappings which do not satisfy Hyers–Ulam stability, Proc. Amer. Math. Soc. 114 (1992), 989–993.
- [13] S. M. Ulam, Problems in Modern Mathematics, Chapter VI, Science Editions, Wiley, New York, 1964.

Choonkil Baak

Department of Mathematics, Chungnam National University, Daejeon 305-764, South Korea
e-mail: cgpark@cnu.ac.kr

Mohammad Sal Moslehian

Department of Mathematics, Ferdowsi University, P. O. Box 1159, Mashhad 91775, Iran
e-mail: moslehian@ferdowsi.um.ac.ir